


Even The TOUGHEST Pathogens cannot RESIST

OFTAQUIX®

Levofloxacin ophthalmic solution 1.5% w/v

Stronger Faster Deeper



* © Iqvia 2019 Midas 2019q2mat In Atc3 S1a Product Ranking (Cravit Family) For Asian Markets: China, Korea, Vietnam, Philippines, Indonesia, Thailand, Taiwan, Malaysia, Singapore, Hong Kong, And India

Disclaimer: Oftaquix and Cravit are brand name of the same product in different countries manufactured by Santen Pharmaceutical Co., Ltd. For the use of registered medical practitioner or a Hospital or a laboratory only.

Abridged information:

INDICATION: Oftaquix 1.5% (Levofloxacin ophthalmic solution 1.5% w/v) is indicated for the treatment of bacterial infections of the eye caused by susceptible microorganisms. **DOSE AND METHOD OF ADMINISTRATION:** Usually, instill one drop a time in the affected eye(s), 3 times per day. **CONTRAINDICATIONS:** Oftaquix 1.5% (Levofloxacin ophthalmic solution 1.5% w/v) is contraindicated in patients with a history of hypersensitivity to the ingredient of this product, ofloxacin or any quinolone antibiotics **USE IN SPECIAL POPULATIONS:** The safety of this product during pregnancy and in low birth weight infants, neonates, infants or children under the age of 15 has not been established. No adjustment of dosage is required in Geriatric population. **WARNING:** Not for injection. For external use only. **PRECAUTION:** In order to avoid the emergence of resistant bacteria, bacterial susceptibility should be confirmed and treatment with Oftaquix 1.5% should be limited to the minimum period required for the eradication of the infection. **ADVERSE REACTIONS:** Shock, anaphylaxis reaction (incidences unknown), but patient should be carefully observed, if reported with shock and anaphylaxis reaction. If any symptoms such as erythema, rash, dyspnoea, decreased blood pressure or eyelid edema etc. are observed then discontinue use and consult physician for appropriate measures. Other adverse reactions (5% >= 0.1%) hypersensitivity (Urticaria, itching), ophthalmic (irritation), others (Dysgeusia (taste bitter etc.)) have been observed. **PACKAGING INFORMATION:** Available in 5 ml plastic dropper bottle **STORAGE AND HANDLING INSTRUCTIONS:** Keep below 30°C, under protection from light. Use the solution within one month after opening the container. For further information, please refer to full pack insert

SANTEN INDIA PRIVATE LIMITED

Unit No - 201 A wing, Sagar Tech Plaza, Saki Naka, Andheri (E), Mumbai – 400072. Phone : 022-49740247
RO : #1101, 3rd floor 24th main, JP Nagar first phase , Bangalore – 560078.